National Oesophago-Gastric Cancer Audit

Dartford and Gravesham NHS Trust

Kent and Medway, Cancer Alliance

Management of High Grade Dysplasia patients

Time period: 2018-22

| | Cancer Alliance | | National |
|---|-----------------|--------|----------|
| Number of High Grade Dysplasia patients in audit | 26 | | 1120 |
| | Cancer Alliance | | National |
| Method of Diagnosis | | | |
| First diagnosis confirmed by second pathologist | N/A | 100.0% | 90.0% |
| Treatment planning | | | |
| – HGD plan discussed at MDT | N/A | 100.0% | 93.0% |
| - Treatment plan for active treatment | N/A | 96.2% | 82.0% |
| First treatment | | | |
| - Endoscopic therapy (such as endoscopic resection) | N/A | 88.5% | 78.0% |
| – Surveillance | N/A | 0.0% | 11.0% |

NOTE: Percentages have been calculated based on the number of patients with recorded data for each variable and not the total number of patients in the Cancer Alliance.

Management of oesophago-gastric cancer patients

Time period: 2020-2022

| | Organisation | | National | |
|---|--------------|-------|----------|--|
| Number of oesophago-gastric (OG) cancer patients in audit | 106 | | 19,865 | |
| Case ascertainment for OG cancer (%) | 85-100% | | | |
| Method of Diagnosis | | | | |
| Patients diagnosed after an emergency admission** ** Percent figure accounts for differences in mix of patients across organisations | 17 | 17.1% | 13.1% | |
| Patients with referral source reported as unknown | 0 | 0.0% | 1.4% | |
| Pre-treatment staging of cancer | | | | |
| Patients having staging CT scan recorded | 94 | 88.7% | 94.0% | |
| Treatment planning | | | | |
| Patients having a plan for curative treatment | 38 | 35.9% | 37.2% | |
| Patients having a plan for non-curative treatment | 68 | 64.2% | 62.8% | |
| - Patients with non-curative plans having chemotherapy or radiotherapy | N/A | | 34.7% | |