

National Oesophago-Gastric Cancer Audit

University Hospitals Dorset NHS Foundation Trust

Wessex, Cancer Alliance

Management of High Grade Dysplasia patients

Time period: 2018-22

		Cancer Alliance	National
Number of High Grade Dysplasia patients in audit	58		1120
		Cancer Alliance	National
Method of Diagnosis			
– First diagnosis confirmed by second pathologist	N/A	90.0%	90.0%
Treatment planning			
– HGD plan discussed at MDT	N/A	93.1%	93.0%
– Treatment plan for active treatment	N/A	82.8%	82.0%
First treatment			
– Endoscopic therapy (such as endoscopic resection)	N/A	81.0%	78.0%
– Surveillance	N/A	12.1%	11.0%


NOTE: Percentages have been calculated based on the number of patients with recorded data for each variable and not the total number of patients in the Cancer Alliance.

Management of oesophago-gastric cancer patients

Time period: 2020-2022


	Organisation		National
Number of oesophago-gastric (OG) cancer patients in audit	227		19,865
Case ascertainment for OG cancer (%)	85-100%		
Method of Diagnosis			
– Patients diagnosed after an emergency admission** ** Percent figure accounts for differences in mix of patients across organisations	41	18.8%	13.1%
– Patients with referral source reported as unknown	0	0.0%	1.4%
Pre-treatment staging of cancer			
– Patients having staging CT scan recorded	198	87.2%	94.0%
Treatment planning			
– Patients having a plan for curative treatment	98	43.2%	37.2%
– Patients having a plan for non-curative treatment	129	56.8%	62.8%
– Patients with non-curative plans having chemotherapy or radiotherapy	40	31.0%	34.7%

90-day mortality rate



Trust	Number	Adjusted %
University Hospitals Dorset NHS Foundation Trust	72	2.75

30-day mortality rate



Trust	Number	Adjusted %
University Hospitals Dorset NHS Foundation Trust	72	0

Outcomes for OG cancer patients who have curative surgery

Time period: 2019-2022

	Organisation	National
Number of patients having surgery	72	5452
Length of stay, median (days)	9	10
Operations in which 15 or more lymph nodes were examined (%)	83.1%	90.1%
30-day postoperative mortality (%)	0.0%	1.5%
90-day postoperative mortality (%)	2.8%	2.9%
Oesophageal surgery		
Number of patients having oesophageal surgery	55	N/A
Oesophageal patients with positive longitudinal margins (%)	14.8%	4.9%
Oesophageal patients with positive circumferential margins (%)	13.7%	21.5%
Gastric surgery		
Number of patients having gastric surgery	17	N/A
Gastric patients with positive longitudinal margins (%)	19.9%	9.8%