

National Oesophago-Gastric Cancer Audit

Mid and South Essex NHS Foundation Trust

East of England - South, Cancer Alliance

Management of High Grade Dysplasia patients

Time period: 2018-22

		Cancer Alliance	National
Number of High Grade Dysplasia patients in audit	72		1120
		Cancer Alliance	National
Method of Diagnosis			
– First diagnosis confirmed by second pathologist	N/A	94.1%	90.0%
Treatment planning			
– HGD plan discussed at MDT	N/A	94.4%	93.0%
– Treatment plan for active treatment	N/A	85.7%	82.0%
First treatment			
– Endoscopic therapy (such as endoscopic resection)	N/A	80.0%	78.0%
– Surveillance	N/A	8.6%	11.0%


NOTE: Percentages have been calculated based on the number of patients with recorded data for each variable and not the total number of patients in the Cancer Alliance.

Management of oesophago-gastric cancer patients

Time period: 2020-2022


	Organisation		National
Number of oesophago-gastric (OG) cancer patients in audit	328		19,865
Case ascertainment for OG cancer (%)	65-74%		
Method of Diagnosis			
– Patients diagnosed after an emergency admission** ** Percent figure accounts for differences in mix of patients across organisations	52	16.8%	13.1%
– Patients with referral source reported as unknown	1	0.3%	1.4%
Pre-treatment staging of cancer			
– Patients having staging CT scan recorded	311	94.8%	94.0%
Treatment planning			
– Patients having a plan for curative treatment	126	38.4%	37.2%
– Patients having a plan for non-curative treatment	202	61.6%	62.8%
– Patients with non-curative plans having chemotherapy or radiotherapy	54	26.7%	34.7%

90-day mortality rate



Trust	Number	Adjusted %
Mid and South Essex NHS Foundation Trust	137	2.57

30-day mortality rate



Trust	Number	Adjusted %
Mid and South Essex NHS Foundation Trust	137	0.67

Outcomes for OG cancer patients who have curative surgery

Time period: 2019-2022

	Organisation	National
Number of patients having surgery	137	5452
Length of stay, median (days)	9	10
Operations in which 15 or more lymph nodes were examined (%)	95.2%	90.1%
30-day postoperative mortality (%)	0.7%	1.5%
90-day postoperative mortality (%)	2.6%	2.9%
Oesophageal surgery		
Number of patients having oesophageal surgery	108	N/A
Oesophageal patients with positive longitudinal margins (%)	19.0%	4.9%
Oesophageal patients with positive circumferential margins (%)	36.0%	21.5%
Gastric surgery		
Number of patients having gastric surgery	29	N/A
Gastric patients with positive longitudinal margins (%)	5.0%	9.8%