

National Oesophago-Gastric Cancer Audit

Norfolk and Norwich University Hospitals NHS Foundation Trust

East of England - North, Cancer Alliance

Management of High Grade Dysplasia patients

Time period: 2018-22

		Cancer Alliance		National
Number of High Grade Dysplasia patients in audit	131			1120
		Cancer Alliance		National
Method of Diagnosis				
– First diagnosis confirmed by second pathologist	N/A	99.2%		90.0%
Treatment planning				
– HGD plan discussed at MDT	N/A	97.0%		93.0%
– Treatment plan for active treatment	N/A	89.3%		82.0%
First treatment				
– Endoscopic therapy (such as endoscopic resection)	N/A	89.3%		78.0%
– Surveillance	N/A	7.6%		11.0%


NOTE: Percentages have been calculated based on the number of patients with recorded data for each variable and not the total number of patients in the Cancer Alliance.

Management of oesophago-gastric cancer patients

Time period: 2020-2022


	Organisation		National
Number of oesophago-gastric (OG) cancer patients in audit	259		19,865
Case ascertainment for OG cancer (%)	85-100%		
Method of Diagnosis			
– Patients diagnosed after an emergency admission** ** Percent figure accounts for differences in mix of patients across organisations	54	21.1%	13.1%
– Patients with referral source reported as unknown	0	0.0%	1.4%
Pre-treatment staging of cancer			
– Patients having staging CT scan recorded	247	95.4%	94.0%
Treatment planning			
– Patients having a plan for curative treatment	85	32.8%	37.2%
– Patients having a plan for non-curative treatment	174	67.2%	62.8%
– Patients with non-curative plans having chemotherapy or radiotherapy	63	36.2%	34.7%

90-day mortality rate



Trust	Number	Adjusted %
Norfolk and Norwich University Hospitals NHS Foundation Trust	135	2.35

30-day mortality rate



Trust	Number	Adjusted %
Norfolk and Norwich University Hospitals NHS Foundation Trust	135	0.65

Outcomes for OG cancer patients who have curative surgery

Time period: 2019-2022

	Organisation	National
Number of patients having surgery	135	5452
Length of stay, median (days)	7	10
Operations in which 15 or more lymph nodes were examined (%)	97.7%	90.1%
30-day postoperative mortality (%)	0.7%	1.5%
90-day postoperative mortality (%)	2.4%	2.9%
Oesophageal surgery		
Number of patients having oesophageal surgery	83	N/A
Oesophageal patients with positive longitudinal margins (%)	4.4%	4.9%
Oesophageal patients with positive circumferential margins (%)	8.6%	21.5%
Gastric surgery		
Number of patients having gastric surgery	52	N/A
Gastric patients with positive longitudinal margins (%)	6.1%	9.8%