

National Oesophago-Gastric Cancer Audit

Bedford Hospital NHS Trust

East of England - South, Cancer Alliance

Management of High Grade Dysplasia patients

Time period: 2015-19

| | | Cancer Alliance | | National |
|---|----|-----------------|-------|----------|
| Number of High Grade Dysplasia patients in audit | 62 | | | 1120 |
| | | Cancer Alliance | | National |
| Method of Diagnosis | | | | |
| – First diagnosis confirmed by second pathologist | 54 | | 87.5% | 90.0% |
| Treatment planning | | | | |
| – HGD plan discussed at MDT | 57 | | 91.9% | 93.0% |
| – Treatment plan for active treatment | 49 | | 79.7% | 82.0% |
| First treatment | | | | |
| – Endoscopic therapy (such as endoscopic resection) | 44 | | 71.2% | 78.0% |
| – Surveillance | 6 | | 10.2% | 11.0% |

NOTE: Percentages have been calculated based on the number of patients with recorded data for each variable and not the total number of patients in the Cancer Alliance.

Management of oesophago-gastric cancer patients

Time period: 2017-2019

| | Organisation | | National |
|---|--------------|-------|----------|
| Number of oesophago-gastric (OG) cancer patients in audit | 75 | | 19,865 |
| Case ascertainment for OG cancer (%) | 75-84% | | |
| Method of Diagnosis | | | |
| – Patients diagnosed after an emergency admission** ** Percent figure accounts for differences in mix of patients across organisations | 4 | 5.8% | 13.1% |
| – Patients with referral source reported as unknown | 0 | 0.0% | 1.4% |
| Pre-treatment staging of cancer | | | |
| – Patients having staging CT scan recorded | 68 | 90.7% | 94.0% |
| Treatment planning | | | |
| – Patients having a plan for curative treatment | 32 | 42.7% | 37.2% |
| – Patients having a plan for non-curative treatment | 43 | 57.3% | 62.8% |
| – Patients with non-curative plans having chemotherapy or radiotherapy | 24 | 55.8% | 34.7% |