

National Oesophago-Gastric Cancer Audit

Leeds Teaching Hospitals NHS Trust

West Yorkshire and Harrogate, Cancer Alliance

Management of High Grade Dysplasia patients

Time period: 2018-22

		Cancer Alliance	National
Number of High Grade Dysplasia patients in audit	18		1120
		Cancer Alliance	National
Method of Diagnosis			
– First diagnosis confirmed by second pathologist	N/A	76.5%	90.0%
Treatment planning			
– HGD plan discussed at MDT	N/A	88.9%	93.0%
– Treatment plan for active treatment	N/A	83.3%	82.0%
First treatment			
– Endoscopic therapy (such as endoscopic resection)	N/A	83.3%	78.0%
– Surveillance	N/A	5.6%	11.0%


NOTE: Percentages have been calculated based on the number of patients with recorded data for each variable and not the total number of patients in the Cancer Alliance.

Management of oesophago-gastric cancer patients

Time period: 2020-2022


	Organisation		National
Number of oesophago-gastric (OG) cancer patients in audit	242		19,865
Case ascertainment for OG cancer (%)	85-100%		
Method of Diagnosis			
– Patients diagnosed after an emergency admission** ** Percent figure accounts for differences in mix of patients across organisations	<10	0.0%	13.1%
– Patients with referral source reported as unknown	86	35.5%	1.4%
Pre-treatment staging of cancer			
– Patients having staging CT scan recorded	238	98.4%	94.0%
Treatment planning			
– Patients having a plan for curative treatment	113	46.7%	37.2%
– Patients having a plan for non-curative treatment	129	53.3%	62.8%
– Patients with non-curative plans having chemotherapy or radiotherapy	N/A		34.7%

90-day mortality rate



Trust	Number	Adjusted %
Leeds Teaching Hospitals NHS Trust	162	2.54

30-day mortality rate



Trust	Number	Adjusted %
Leeds Teaching Hospitals NHS Trust	162	1.86

Outcomes for OG cancer patients who have curative surgery

Time period: 2019-2022

	Organisation	National
Number of patients having surgery	162	5452
Length of stay, median (days)	10	10
Operations in which 15 or more lymph nodes were examined (%)	87.6%	90.1%
30-day postoperative mortality (%)	1.9%	1.5%
90-day postoperative mortality (%)	2.5%	2.9%
Oesophageal surgery		
Number of patients having oesophageal surgery	112	N/A
Oesophageal patients with positive longitudinal margins (%)	11.8%	4.9%
Oesophageal patients with positive circumferential margins (%)	29.8%	21.5%
Gastric surgery		
Number of patients having gastric surgery	50	N/A
Gastric patients with positive longitudinal margins (%)	17.5%	9.8%