National Oesophago-Gastric Cancer Audit

The Royal Marsden NHS Foundation Trust **

RM Partners West London, Cancer Alliance

Management of High Grade Dysplasia patients

Time period: 2018-22

	Cancer Alliance		National
Number of High Grade Dysplasia patients in audit	48		1120
	Cancer Alliance		National
Method of Diagnosis			
- First diagnosis confirmed by second pathologist	N/A	95.5%	90.0%
Treatment planning			
- HGD plan discussed at MDT	N/A	95.8%	93.0%
- Treatment plan for active treatment	N/A	85.0%	82.0%
First treatment			
- Endoscopic therapy (such as endoscopic resection)	N/A	85.0%	78.0%
- Surveillance	N/A	2.5%	11.0%


NOTE: Percentages have been calculated based on the number of patients with recorded data for each variable and not the total number of patients in the Cancer Alliance.

Management of oesophago-gastric cancer patients

Time period: 2020-2022


	Organisation	National	
Number of oesophago-gastric (OG) cancer patients in audit	N/A	19,865	
Case ascertainment for OG cancer (%)			
Method of Diagnosis			
 Patients diagnosed after an emergency admission** ** Percent figure accounts for differences in mix of patients across organisations 	N/A	13.1%	
Patients with referral source reported as unknown	N/A	1.4%	
Pre-treatment staging of cancer			
- Patients having staging CT scan recorded	N/A	94.0%	
Treatment planning			
- Patients having a plan for curative treatment	N/A	37.2%	
- Patients having a plan for non-curative treatment	N/A	62.8%	
- Patients with non-curative plans having chemotherapy or radiotherapy	N/A	34.7%	

90-day mortality rate



Trust	Number	Adjusted %
The Royal Marsden NHS Foundation Trust **	154	3.25

30-day mortality rate



Trust	Number	Adjusted %
The Royal Marsden NHS Foundation Trust **	154	2.53

Outcomes for OG cancer patients who have curative surgery

Time period: 2019-2022

	Organisation	National
Number of patients having surgery	154	5452
Length of stay, median (days)	12	10
Operations in which 15 or more lymph nodes were examined (%)	97.4%	90.1%
30-day postoperative mortality (%)	2.5%	1.5%
90-day postoperative mortality (%)	3.3%	2.9%
Oesophageal surgery		
Number of patients having oesophageal surgery	110	N/A
Oesophageal patients with positive longitudinal margins (%)	6.1%	4.9%
Oesophageal patients with positive circumferential margins (%)	11.7%	21.5%
Gastric surgery		
Number of patients having gastric surgery	44	N/A
Gastric patients with positive longitudinal margins (%)	11.9%	9.8%