National Oesophago-Gastric Cancer Audit

Liverpool University Hospitals NHS Foundation Trust

Cheshire and Merseyside, Cancer Alliance

Management of High Grade Dysplasia patients

Time period: 2018-22

	Cancer Alliance		National
Number of High Grade Dysplasia patients in audit	13		1120
	Cancer Alliance		National
Method of Diagnosis			
- First diagnosis confirmed by second pathologist	N/A	100.0%	90.0%
Treatment planning			
- HGD plan discussed at MDT	N/A	100.0%	93.0%
- Treatment plan for active treatment	N/A	100.0%	82.0%
First treatment			
- Endoscopic therapy (such as endoscopic resection)	N/A	85.7%	78.0%
- Surveillance	N/A	0.0%	11.0%


NOTE: Percentages have been calculated based on the number of patients with recorded data for each variable and not the total number of patients in the Cancer Alliance.

Management of oesophago-gastric cancer patients

Time period: 2020-2022


	Organisation		National	
Number of oesophago-gastric (OG) cancer patients in audit	294		19,865	
Case ascertainment for OG cancer (%)	85-100%			
Method of Diagnosis				
 Patients diagnosed after an emergency admission** ** Percent figure accounts for differences in mix of patients across organisations 	33	10.3%	13.1%	
Patients with referral source reported as unknown	0	0.0%	1.4%	
Pre-treatment staging of cancer				
- Patients having staging CT scan recorded	N/A		94.0%	
Treatment planning				
- Patients having a plan for curative treatment	118	40.1%	37.2%	
- Patients having a plan for non-curative treatment	176	59.9%	62.8%	
- Patients with non-curative plans having chemotherapy or radiotherapy	57	32.4%	34.7%	

90-day mortality rate



Trust	Number	Adjusted %
Liverpool University Hospitals NHS Foundation Trust	82	2.1

30-day mortality rate



Trust	Number	Adjusted %
Liverpool University Hospitals NHS Foundation Trust	82	0

Outcomes for OG cancer patients who have curative surgery

Time period: 2019-2022

	Organisation	National
Number of patients having surgery	82	5452
Length of stay, median (days)	11	10
Operations in which 15 or more lymph nodes were examined (%)	95.9%	90.1%
30-day postoperative mortality (%)	0.0%	1.5%
90-day postoperative mortality (%)	2.1%	2.9%
Oesophageal surgery		
Number of patients having oesophageal surgery	50	N/A
Oesophageal patients with positive longitudinal margins (%)	3.4%	4.9%
Oesophageal patients with positive circumferential margins (%)	17.2%	21.5%
Gastric surgery		
Number of patients having gastric surgery	32	N/A
Gastric patients with positive longitudinal margins (%)	12.9%	9.8%