

National Oesophago-Gastric Cancer Audit

Poole Hospital NHS Foundation Trust

Wessex, Cancer Alliance

Management of High Grade Dysplasia patients

Time period: 2017-21

		Cancer Alliance	National
Number of High Grade Dysplasia patients in audit	57		1120
		Cancer Alliance	National
Method of Diagnosis			
– First diagnosis confirmed by second pathologist	27	90.0%	90.0%
Treatment planning			
– HGD plan discussed at MDT	48	84.2%	93.0%
– Treatment plan for active treatment	44	77.2%	82.0%
First treatment			
– Endoscopic therapy (such as endoscopic resection)	43	97.7%	78.0%
– Surveillance	9	15.8%	11.0%

NOTE: Percentages have been calculated based on the number of patients with recorded data for each variable and not the total number of patients in the Cancer Alliance.

Management of oesophago-gastric cancer patients

Time period: 2018-2020

	Organisation		National
Number of oesophago-gastric (OG) cancer patients in audit	110		19,865
Case ascertainment for OG cancer (%)	85-100%		
Method of Diagnosis			
– Patients diagnosed after an emergency admission** ** Percent figure accounts for differences in mix of patients across organisations	3	3.0%	13.1%
– Patients with referral source reported as unknown	0	0.0%	1.4%
Pre-treatment staging of cancer			
– Patients having staging CT scan recorded	103	93.6%	94.0%
Treatment planning			
– Patients having a plan for curative treatment	44	40.0%	37.2%
– Patients having a plan for non-curative treatment	66	60.0%	62.8%
– Patients with non-curative plans having chemotherapy or radiotherapy	37	56.1%	34.7%