

National Oesophago-Gastric Cancer Audit

Basildon and Thurrock University Hospitals NHS Foundation Trust

East of England - South, Cancer Alliance

Management of High Grade Dysplasia patients

Time period: 2015-19

		Cancer Alliance	National
Number of High Grade Dysplasia patients in audit	62		1120
		Cancer Alliance	National
Method of Diagnosis			
– First diagnosis confirmed by second pathologist	54	87.5%	90.0%
Treatment planning			
– HGD plan discussed at MDT	57	91.9%	93.0%
– Treatment plan for active treatment	49	79.7%	82.0%
First treatment			
– Endoscopic therapy (such as endoscopic resection)	44	71.2%	78.0%
– Surveillance	6	10.2%	11.0%

NOTE: Percentages have been calculated based on the number of patients with recorded data for each variable and not the total number of patients in the Cancer Alliance.

Management of oesophago-gastric cancer patients

Time period: 2017-2019

	Organisation		National
Number of oesophago-gastric (OG) cancer patients in audit	55		19,865
Case ascertainment for OG cancer (%)	<65%		
Method of Diagnosis			
– Patients diagnosed after an emergency admission** ** Percent figure accounts for differences in mix of patients across organisations	9	16.8%	13.1%
– Patients with referral source reported as unknown	1	1.8%	1.4%
Pre-treatment staging of cancer			
– Patients having staging CT scan recorded	50	90.9%	94.0%
Treatment planning			
– Patients having a plan for curative treatment	23	41.8%	37.2%
– Patients having a plan for non-curative treatment	32	58.2%	62.8%
– Patients with non-curative plans having chemotherapy or radiotherapy	11	34.4%	34.7%