# **National Oesophago-Gastric Cancer Audit**

## **Swansea Bay University Health Board**

Swansea Bay, Welsh Region

#### **Management of High Grade Dysplasia patients**

Time period: 2018-22

	Welsh Region	National	
Number of High Grade Dysplasia patients in audit	N/A	1120	
	Welsh Region	National	
Method of Diagnosis			
- First diagnosis confirmed by second pathologist	N/A	90.0%	
Treatment planning			
- HGD plan discussed at MDT	N/A	93.0%	
- Treatment plan for active treatment	N/A	82.0%	
First treatment			
- Endoscopic therapy (such as endoscopic resection)	N/A	78.0%	
- Surveillance	N/A	11.0%	


NOTE: Percentages have been calculated based on the number of patients with recorded data for each variable and not the total number of patients in the Cancer Alliance.

### Management of oesophago-gastric cancer patients

Time period: 2020-2022


	Organisation		National
Number of oesophago-gastric (OG) cancer patients in audit	192		19,865
Case ascertainment for OG cancer (%)	85-100%		
Method of Diagnosis			
- Patients diagnosed after an emergency admission**  ** Percent figure accounts for differences in mix of patients across organisations	39	20.5%	13.1%
Patients with referral source reported as unknown	0	0.0%	1.4%
Pre-treatment staging of cancer			
- Patients having staging CT scan recorded	186	96.9%	94.0%
Treatment planning			
- Patients having a plan for curative treatment	49	25.5%	37.2%
- Patients having a plan for non-curative treatment	143	74.5%	62.8%
- Patients with non-curative plans having chemotherapy or radiotherapy	59	41.3%	34.7%

#### 90-day mortality rate



Trust	Number	Adjusted %
Swansea Bay University Health Board	20	0

#### 30-day mortality rate



Trust	Number	Adjusted %
Swansea Bay University Health Board	20	0

### Outcomes for OG cancer patients who have curative surgery

Time period: 2019-2022

	Organisation	National	
Number of patients having surgery	11	5452	
Length of stay, median (days)	N/A	10	
Operations in which 15 or more lymph nodes were examined (%)		90.1%	
30-day postoperative mortality (%)		1.5%	
90-day postoperative mortality (%)		2.9%	
Oesophageal surgery			
Number of patients having oesophageal surgery	N/A	N/A	
Oesophageal patients with positive longitudinal margins (%)		4.9%	
Oesophageal patients with positive circumferential margins (%)		21.5%	
Gastric surgery			
Number of patients having gastric surgery	N/A	N/A	
Gastric patients with positive longitudinal margins (%)		9.8%	