

# National Oesophago-Gastric Cancer Audit

## Nottingham University Hospitals NHS Trust

East Midlands, Cancer Alliance

### Management of High Grade Dysplasia patients

Time period: 2018-22

		Cancer Alliance	National
Number of High Grade Dysplasia patients in audit	120		1120
		Cancer Alliance	National
Method of Diagnosis			
– First diagnosis confirmed by second pathologist	N/A	75.3%	90.0%
Treatment planning			
– HGD plan discussed at MDT	N/A	91.7%	93.0%
– Treatment plan for active treatment	N/A	84.2%	82.0%
First treatment			
– Endoscopic therapy (such as endoscopic resection)	N/A	78.1%	78.0%
– Surveillance	N/A	8.8%	11.0%


NOTE: Percentages have been calculated based on the number of patients with recorded data for each variable and not the total number of patients in the Cancer Alliance.

## Management of oesophago-gastric cancer patients

Time period: 2020-2022


	Organisation		National
Number of oesophago-gastric (OG) cancer patients in audit	317		19,865
Case ascertainment for OG cancer (%)	85-100%		
<b>Method of Diagnosis</b>			
– Patients diagnosed after an emergency admission** ** Percent figure accounts for differences in mix of patients across organisations	11	3.6%	13.1%
– Patients with referral source reported as unknown	1	0.3%	1.4%
<b>Pre-treatment staging of cancer</b>			
– Patients having staging CT scan recorded	304	95.9%	94.0%
<b>Treatment planning</b>			
– Patients having a plan for curative treatment	154	48.6%	37.2%
– Patients having a plan for non-curative treatment	163	51.4%	62.8%
– Patients with non-curative plans having chemotherapy or radiotherapy	65	39.9%	34.7%

90-day mortality rate



Trust	Number	Adjusted %
Nottingham University Hospitals NHS Trust	280	4

30-day mortality rate



Trust	Number	Adjusted %
Nottingham University Hospitals NHS Trust	280	2.08

## Outcomes for OG cancer patients who have curative surgery

Time period: 2019-2022

	Organisation	National
Number of patients having surgery	280	5452
Length of stay, median (days)	9	10
Operations in which 15 or more lymph nodes were examined (%)	77.4%	90.1%
30-day postoperative mortality (%)	2.1%	1.5%
90-day postoperative mortality (%)	4.0%	2.9%
<b>Oesophageal surgery</b>		
Number of patients having oesophageal surgery	201	N/A
Oesophageal patients with positive longitudinal margins (%)	8.5%	4.9%
Oesophageal patients with positive circumferential margins (%)	28.4%	21.5%
<b>Gastric surgery</b>		
Number of patients having gastric surgery	79	N/A
Gastric patients with positive longitudinal margins (%)	18.9%	9.8%