

National Oesophago-Gastric Cancer Audit

Royal Surrey County Hospital NHS Foundation Trust

Surrey and Sussex, Cancer Alliance

Management of High Grade Dysplasia patients

Time period: 2018-22

		Cancer Alliance		National
Number of High Grade Dysplasia patients in audit	37			1120
		Cancer Alliance		National
Method of Diagnosis				
– First diagnosis confirmed by second pathologist	N/A	90.3%	90.0%	
Treatment planning				
– HGD plan discussed at MDT	N/A	94.6%	93.0%	
– Treatment plan for active treatment	N/A	81.3%	82.0%	
First treatment				
– Endoscopic therapy (such as endoscopic resection)	N/A	65.6%	78.0%	
– Surveillance	N/A	6.3%	11.0%	


NOTE: Percentages have been calculated based on the number of patients with recorded data for each variable and not the total number of patients in the Cancer Alliance.

Management of oesophago-gastric cancer patients

Time period: 2020-2022


	Organisation		National
Number of oesophago-gastric (OG) cancer patients in audit	115		19,865
Case ascertainment for OG cancer (%)	75-84%		
Method of Diagnosis			
– Patients diagnosed after an emergency admission** ** Percent figure accounts for differences in mix of patients across organisations	<10	8.1%	13.1%
– Patients with referral source reported as unknown	0	0.0%	1.4%
Pre-treatment staging of cancer			
– Patients having staging CT scan recorded	110	95.7%	94.0%
Treatment planning			
– Patients having a plan for curative treatment	52	45.2%	37.2%
– Patients having a plan for non-curative treatment	63	54.8%	62.8%
– Patients with non-curative plans having chemotherapy or radiotherapy	N/A		34.7%

90-day mortality rate



Trust	Number	Adjusted %
Royal Surrey County Hospital NHS Foundation Trust	128	3.11

30-day mortality rate



Trust	Number	Adjusted %
Royal Surrey County Hospital NHS Foundation Trust	128	1.65

Outcomes for OG cancer patients who have curative surgery

Time period: 2019-2022

	Organisation	National
Number of patients having surgery	128	5452
Length of stay, median (days)	12	10
Operations in which 15 or more lymph nodes were examined (%)	100.0%	90.1%
30-day postoperative mortality (%)	1.7%	1.5%
90-day postoperative mortality (%)	3.1%	2.9%
Oesophageal surgery		
Number of patients having oesophageal surgery	87	N/A
Oesophageal patients with positive longitudinal margins (%)	2.1%	4.9%
Oesophageal patients with positive circumferential margins (%)	2.8%	21.5%
Gastric surgery		
Number of patients having gastric surgery	41	N/A
Gastric patients with positive longitudinal margins (%)	0.0%	9.8%